

# Elliptic curves over quadratic fields with fixed torsion subgroup and positive rank

Mirela Jukić Bokun

## Abstract

For each of the torsion groups  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/10\mathbb{Z}$ ,  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/12\mathbb{Z}$ ,  $\mathbb{Z}/15\mathbb{Z}$  we find the quadratic field with the smallest absolute value of its discriminant such that there exists an elliptic curve with that torsion and positive rank. For the torsion groups  $\mathbb{Z}/11\mathbb{Z}$ ,  $\mathbb{Z}/14\mathbb{Z}$  we solve the analogous problem after assuming the Parity conjecture.

**Keywords:** elliptic curve, torsion group, rank, quadratic field

**2000 Mathematics Subject Classification:** 11G05, 14H52, 11R11

## 1 Introduction

According to the Mordell-Weil theorem, the group of  $K$ -rational points  $E(K)$  of an elliptic curve  $E$  over a number field  $K$  is a finitely generated abelian group. Therefore, the group  $E(K)$  is isomorphic to  $E(K)_{\text{tors}} \oplus \mathbb{Z}^r$ , where  $E(K)_{\text{tors}}$  is the torsion subgroup and the non-negative integer  $r$  is the rank. When  $K = \mathbb{Q}$ , by Mazur's theorem [11] the torsion group is one of the following 15 groups:  $\mathbb{Z}/n\mathbb{Z}$  with  $1 \leq n \leq 10$  or  $n = 12$ ,  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2m\mathbb{Z}$  with  $1 \leq m \leq 4$ . If  $K$  is a quadratic field, the following theorem of Kenku and Momose [9] and Kamienny [7] describes possible torsions.

**Theorem 1** *Let  $K$  be a quadratic field and  $E$  an elliptic curve over  $K$ . Then the torsion subgroup  $E(K)_{\text{tors}}$  of  $E(K)$  is isomorphic to one of the following 26 groups:*

- $\mathbb{Z}/n\mathbb{Z}$ , for  $1 \leq n \leq 16$ ,  $n = 18$ ,
- $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2n\mathbb{Z}$ , for  $1 \leq n \leq 6$ ,

---

<sup>0</sup>The author was supported by the Ministry of Science, Education and Sports, Republic of Croatia, grant 235-2352818-1042

- $\mathbb{Z}/3\mathbb{Z} \oplus \mathbb{Z}/3n\mathbb{Z}$ , for  $n = 1, 2$ ,  $K = \mathbb{Q}(\sqrt{-3})$ ,
- $\mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$  for  $K = \mathbb{Q}(i)$ .

Kamienny and Najman [8] described methods that can be used to find all possible torsions over a given quadratic field. For each possible torsion group, they found the exact field with the smallest absolute value of the discriminant  $\Delta$  such that an elliptic curve with that torsion group exists. Curves with positive ranks over quadratic fields can be found in [2, 5, 6, 12]. In [2], the authors give examples of elliptic curves with given torsion and positive rank over fields of degree 2, 3 and 4.

In this article we use techniques from [8] in order to establish, for each of torsion groups  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/10\mathbb{Z}$ ,  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/12\mathbb{Z}$ ,  $\mathbb{Z}/15\mathbb{Z}$ , the quadratic field with the smallest absolute value of the discriminant such that there exists an elliptic curve with that torsion and positive rank. In the case of the torsion subgroups  $\mathbb{Z}/11\mathbb{Z}$ ,  $\mathbb{Z}/14\mathbb{Z}$  the analogous problem is solved conditionally - we conclude that, if the Parity conjecture holds (i.e. the algebraic rank of the elliptic curve  $E$  over a number field  $K$  has the same parity as the analytic rank which is given by the root number (see, for example, [4]), then the elliptic curves that we find have positive rank.

We performed a search in infinite families of elliptic curves with given torsion to find elliptic curves with positive rank over the quadratic field with the smallest absolute value of the discriminant. All torsion groups that we deal with have the common property that they come from families of elliptic curves with prescribed torsion which are parameterized by another elliptic curve.

For rank computation we use Cremona's MWRANK [3] (for elliptic curve with rational coefficients) and an implementation of 2-descent over number fields [13] of Simon in PARI/GP. We also use MAGMA [1] for computing 2-Selmer ranks and root numbers.

## 2 Elliptic curves with given torsions and positive ranks over smallest fields

### 2.1 Torsion group $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/10\mathbb{Z}$

First we will derive an equation for elliptic curve with torsion subgroup  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/10\mathbb{Z}$ .

According to [10], the elliptic curve with equation

$$\begin{aligned}\mathcal{E}^{(10)} &: y^2 + \frac{2t^3 + 2t^2 + 2t + 1}{(t+1)^2}xy + \frac{t^2(2t+1)}{(t+1)^2}y \\ &= x^3 + \frac{t^2(2t+1)}{(t+1)^2}x^2\end{aligned}$$

has a  $\mathbb{Q}(t)$ -rational torsion point

$$Q = \left( -\frac{t^2(2t+1)}{(t+1)^3}, -\frac{t^3(2t+1)^2}{(t+1)^5} \right)$$

of order 10. It can be shown that the curve  $\mathcal{E}^{(10)}$  is birationally equivalent to the elliptic curve

$$\begin{aligned}\mathcal{E}'^{(10)} &: y^2 = x^3 - 2(1+2t-5t^2-5t^4-2t^5+t^6)x^2 \\ &\quad + (-1+t^2)^5(-1-4t+t^2)x.\end{aligned}\tag{1}$$

The right hand side of the equation (1) has a linear and a quadratic factor. The curve will have more points of order 2 over the field  $K$  if the discriminant of the quadratic factor is a full square in  $K$ . This leads to condition that if the parameter  $t$  is  $x$ -coordinate of point on the curve  $X_1(2, 10)(K)$ , where

$$X_1(2, 10): s^2 = t^3 + t^2 - t,$$

then the curve  $\mathcal{E}'^{(10)}(K)$  has torsion subgroup  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/10\mathbb{Z}$  generated by the points

$$\begin{aligned}T_2 &= (1+2t-5t^2-5t^4-2t^5+t^6-8t^2s, 0), \\ T_{10} &= ((1+t)^3(1+3t-5t^2+t^3), -4t(1+t)^3(1+3t-5t^2+t^3)).\end{aligned}$$

**Theorem 2** *The quadratic field  $K$  with smallest  $|\Delta|$  over which there appears an elliptic curve with a torsion group  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/10\mathbb{Z}$  and positive rank is  $K = \mathbb{Q}(\sqrt{-2})$ .*

PROOF. Note that, according to [8], the minimal quadratic field  $K$  such that  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/10\mathbb{Z}$  appears as a torsion group over  $K$  is  $K = \mathbb{Q}(\sqrt{-2})$ . The elliptic curve  $X_1(2, 10)$  over the field  $\mathbb{Q}(\sqrt{-2})$  has rank equal to 1 with the generator  $P = (-2, -\sqrt{-2})$ . The elliptic curve generated from (1) by the point  $2P$ , i.e. by the parameters

$$(t, s) = \left( -\frac{25}{8}, \frac{95}{32}\sqrt{-2} \right),$$

is

$$E^{(2,10)} : y^2 = x^3 - \frac{261214369}{131072}x^2 + \frac{75626226572068161}{68719476736}x.$$

This elliptic curve has rational coefficients and we can use MWRANK to show that it has rank equal to 1 over  $\mathbb{Q}$ . The rank of this curve over  $\mathbb{Q}(\sqrt{-2})$  is then at least 1 and that proves the theorem.  $\square$

**Remark 1** *It can be shown that the elliptic curve from the last proof has rank equal to 3 over the field  $K = \mathbb{Q}(\sqrt{-2})$  (using the fact that if  $E$  is an elliptic curve over  $\mathbb{Q}$ , then the rank of  $E$  over  $\mathbb{Q}(\sqrt{-2})$  is given by  $\text{rank}(E(\mathbb{Q}(\sqrt{-2}))) = \text{rank}(E(\mathbb{Q})) + \text{rank}(E_{-2}(\mathbb{Q}))$ , where  $E_{-2}$  is the  $(-2)$ -twist of  $E$  over  $\mathbb{Q}$ ).*

## 2.2 Torsion group $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/12\mathbb{Z}$

According to [10], the point

$$Q = \left( -\frac{1}{8}(t+1)(t^2 - 2t + 5), -\frac{1}{32}(t+1)(t^2 - 1)(t^2 - 2t + 5) \right)$$

is a torsion point of the order 12 of the elliptic curve

$$\mathcal{E}^{(12)} : y^2 - \frac{(t^4 - 6t^2 + 24t - 3)}{4(t-1)^2}xy - \frac{(t+1)^2(t^2 - 2t + 5)}{4(t-1)^2}y = x^3.$$

It can be shown that this curve is birationally equivalent to the curve

$$\begin{aligned} \mathcal{E}^{(2,12)} : y^2 &= x^3 + 256(1 - 4t + 4t^2 + 20t^3 - 26t^4 + 20t^5 + 4t^6 - 4t^7 + t^8) \\ &\quad \times (1 + 4t + 4t^3 + t^4 - 4z - 2t^2(1 + 2z))x^2 \\ &\quad + 1048576(t-1)^6(t+1)^2(t-z)^6(-1 + t - t^2 + z)^2x. \end{aligned} \quad (2)$$

Similarly to the previous case, this curve has more points of order 2 over the field  $K$  if  $(t, z) \in X_1(2, 12)(K)$ , where

$$X_1(2, 12) : z^2 = t^3 - t^2 + t.$$

In that case, points

$$\begin{aligned} T_2 &= (-4096t^3(t^2 - t + 1)(1 + 4t + 4t^3 + t^4 - 4z - 2t^2(1 + 2z)), 0), \\ T_{12} &= (-1024(t-1)^5(t+1)(-2t - 2t^3 + z + t^2(2+z)), \\ &\quad -16384(t-1)^5(1+t+t^2+t^3)(-6t^7 - z + t^6(-2+z) - 5t^2(2+z) \\ &\quad + 6t^5(1+2z) + t(2+4z) + 2t^3(7+8z) - t^4(20+11z))). \end{aligned}$$

generate a torsion subgroup  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/12\mathbb{Z}$  on  $\mathcal{E}^{(2,12)}(K)$ .

**Theorem 3** *The quadratic field  $K$  with smallest  $|\Delta|$  over which there appears an elliptic curve with a torsion group  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/12\mathbb{Z}$  and positive rank is  $K = \mathbb{Q}(\sqrt{13})$ .*

PROOF. According to [8], the minimal quadratic field  $K$  such that  $\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/12\mathbb{Z}$  appears as a torsion group over  $K$  is  $K = \mathbb{Q}(\sqrt{13})$ . The elliptic curve  $X_1(2, 12)$  over the field  $\mathbb{Q}(\sqrt{13})$  has rank equal to 1 with the generator  $P = (-2, -\sqrt{-2})$ . The elliptic curve generated from (2) by point  $P$  is

$$E^{(2,12)} : \quad y^2 = x^3 + \frac{1}{65536}(-4289032\sqrt{13} + 15673889)x^2 \\ + \frac{1}{4194304}(-32028469200\sqrt{13} + 115490749725)x,$$

and has rank equal to 1. A point of infinite order is

$$Q = \left( -\frac{153}{256}\sqrt{13} + \frac{4473}{2048}, -\frac{9046125}{131072}\sqrt{13} + \frac{130609215}{524288} \right).$$

□

### 2.3 Torsion group $\mathbb{Z}/15\mathbb{Z}$

According to [12], the general equation of the elliptic curve with points of order 15 is

$$\mathcal{E}^{(15)} : \quad y^2 + axy + by = x^3 + bx^2, \\ a = \frac{(t^2 - t)s + (t^5 + 5t^4 + 9t^3 + 7t^2 + 4t + 1)}{(t + 1)^3(t^2 + t + 1)}, \quad (3) \\ b = \frac{t(t^4 - 2t^2 - t - 1)s + t^3(t + 1)(t^3 + 3t^2 + t + 1)}{(t + 1)^6(t^2 + t + 1)},$$

with  $(t, s) \in X_1(15)$ , where

$$X_1(15) : \quad s^2 + st + s = t^3 + t^2,$$

and

$$t(t + 1)(t^2 + t + 1)(t^4 + 3t^3 + 4t^2 + 2t + 1)(t^4 - 7t^3 - 6t^2 + 2t + 1) \neq 0.$$

**Theorem 4** *The quadratic field  $K$  with smallest  $|\Delta|$  over which there appears an elliptic curve with a torsion group  $\mathbb{Z}/15\mathbb{Z}$  and positive rank is  $K = \mathbb{Q}(\sqrt{-7})$ .*

PROOF. According to [8], the minimal quadratic field  $K$  such that  $\mathbb{Z}/15\mathbb{Z}$  appears as a torsion group over  $K$  is  $K = \mathbb{Q}(\sqrt{5})$  and over this field there is only one elliptic curve with torsion  $\mathbb{Z}/15\mathbb{Z}$  and that curve has rank equal to 0.

Next we look at elliptic curves over the field  $\mathbb{Q}(\sqrt{-7})$ . We find that  $X_1(15)$  over this field has rank equal to 1 with the point

$$P = \left( \frac{1}{8}(-\sqrt{-7} - 5), \frac{1}{16}(\sqrt{-7} - 11) \right)$$

of infinite order. The elliptic curve generated from (3) by the point  $-P$  i.e. point with parameters

$$(t, s) = -P = \left( \frac{1}{8}(-\sqrt{-7} - 5), \frac{1}{16}(\sqrt{-7} - 11) \right) \in X_1(15)(\mathbb{Q}(\sqrt{-7})),$$

is

$$E^{(15)} : y^2 + (-2\sqrt{-7} + 15)xy + (26\sqrt{-7} - 14)y = x^3 + (26\sqrt{-7} - 14)x^2.$$

According to 2-descent procedure implemented in `e11.gp`, this curve has rank  $\leq 1$  over  $\mathbb{Q}(\sqrt{-7})$ . The rootnumber calculated by MAGMA and the Parity conjecture suggest that this curve, conditionally, has rank 1. Finally, in the set

$$\{a + b\sqrt{-7} : |a|, |b| \leq 100\}$$

we successfully find  $x$ -coordinate of the non-torsion point

$$Q = (6\sqrt{-7} - 98, 136\sqrt{-7} + 1064)$$

on the curve  $E^{(15)}(\mathbb{Q}(\sqrt{-7}))$  and this completes the proof.  $\square$

## 2.4 Torsion group $\mathbb{Z}/11\mathbb{Z}$

According to [12], the general equation of the elliptic curve with points of order 11 is

$$\mathcal{E}^{(11)} : \begin{aligned} y^2 + (st + t - s^2)xy + s(s-1)(s-t)t^2y \\ x^3 + s(s-1)(s-t)tx^2, \end{aligned} \quad (4)$$

with  $P = (t, s) \in X_1(11)$ , where

$$X_1(11) : s^2 - s = t^3 - t^2$$

and

$$\Delta_{\mathcal{E}(11)} = t(t-1)(t^5 - 18t^4 + 35t^3 - 16t^2 - 2t + 1) \neq 0.$$

**Theorem 5** *Assume the Parity conjecture. The quadratic field  $K$  with smallest  $|\Delta|$  over which there appears an elliptic curve with a torsion group  $\mathbb{Z}/11\mathbb{Z}$  and positive rank is  $K = \mathbb{Q}(\sqrt{-7})$ .*

PROOF. According to [8], the minimal quadratic field  $K$  such that  $\mathbb{Z}/11\mathbb{Z}$  appears as a torsion group over  $K$  is  $K = \mathbb{Q}(\sqrt{-7})$ . The elliptic curve  $X_1(11)$  over the field  $\mathbb{Q}(\sqrt{-7})$  has rank equal to 1. The point of infinite order is

$$P = \left( -\frac{4}{7}, \frac{1}{98}(49 + 19\sqrt{-7}) \right),$$

and generates, from (4), the elliptic curve

$$\begin{aligned} E^{(11)} : \quad y^2 + (-209\sqrt{-7} - 579)xy - (10486784\sqrt{-7} + 57953280)y \\ = x^3 + (26752\sqrt{-7} + 147840)x^2, \end{aligned}$$

with the torsion group  $\mathbb{Z}/11\mathbb{Z}$ . The root number of this elliptic curve over the field  $\mathbb{Q}(\sqrt{-7})$  is -1 so, according to the Parity conjecture, we can conclude that this is elliptic curve with a conditionally positive rank. Unfortunately, we were unable to determine any point of infinite order on this curve.  $\square$

## 2.5 Torsion group $\mathbb{Z}/14\mathbb{Z}$

The general equation of the elliptic curve with points of order 14, according to [12], is

$$\begin{aligned} \mathcal{E}^{(14)} : \quad y^2 + axy + by = x^3 + bx^2, \\ a = \frac{t^4 - st^3 + (2s - 4)t^2 - st + 1}{(t + 1)(t^3 - 2t^2 - t + 1)}, \\ b = \frac{-t^7 + 2t^6 + (2s - 1)t^5 + (-2s - 1)t^4 + (-2s + 2)t^3 + (3s - 1)t^2 - st}{(t + 1)^2(t^3 - 2t^2 - t + 1)^2}, \end{aligned} \tag{5}$$

where  $P = (t, s) \in X_1(14)$  and

$$\begin{aligned} X_1(14) : \quad s^2 + st + s = t^3 - t, \\ \Delta_{\mathcal{E}(14)} = t(t-1)(t+1)(t^3 - 9t^2 - t + 1)(t^3 - 2t^2 - t + 1) \neq 0. \end{aligned}$$

**Theorem 6** *Assume the Parity conjecture. The quadratic field  $K$  with smallest  $|\Delta|$  over which there appears an elliptic curve with a torsion group  $\mathbb{Z}/14\mathbb{Z}$  and positive rank is  $K = \mathbb{Q}(\sqrt{-11})$ .*

PROOF. According to [8], the minimal quadratic field  $K$  such that  $\mathbb{Z}/14\mathbb{Z}$  appears as a torsion group over  $K$  is  $K = \mathbb{Q}(\sqrt{-7})$ . The elliptic curve  $X_1(14)$  over the field  $\mathbb{Q}(\sqrt{-7})$  has rank equal to 0. It can be shown that the torsion points of this curve which are not the cusps generate two non-isomorphic elliptic curves. Over the fields  $\mathbb{Q}(\sqrt{-2})$  and  $\mathbb{Q}(\sqrt{2})$ , the elliptic curve  $X_1(14)$  has rank equal to 0 and torsion  $\mathbb{Z}/6\mathbb{Z}$ . All the torsion points are cusps so we conclude that elliptic curve with the torsion group  $\mathbb{Z}/14\mathbb{Z}$  over these two fields does not exist.

Over the field  $\mathbb{Q}(\sqrt{-11})$  the curve  $X_1(14)$  has rank equal to 1, with the generator

$$P = \left( -\frac{5}{4}, -\frac{1}{4}\sqrt{-11} + \frac{1}{8} \right).$$

The linear combination  $-2P + T$ , where  $T = (1, -2)$  is a torsion point on  $X_1(14)(\mathbb{Q}(\sqrt{-11}))$ , generates the elliptic curve

$$\begin{aligned} E^{(14)} : & y^2 + (-2601888534886283704\sqrt{-11} + 154252733407512581857)xy \\ & + (-43286504429925775681153399339981518914403432663035092419000\sqrt{-11} \\ & + 239448337641930912934754848966237361950084593095601008052000)y \\ & = x^3 + (-313766195076761969526071169866614175160\sqrt{-11} \\ & + 1735663223649526033628839600302712469280)x^2 \end{aligned}$$

with the torsion group  $\mathbb{Z}/11\mathbb{Z}$ . The root number of this elliptic curve over the field  $\mathbb{Q}(\sqrt{-11})$  is -1 so, if we assume that the Parity conjecture holds, we can conclude that this is the elliptic curve with a conditionally positive rank.  $\square$

## References

- [1] W. BOSMA, J. CANNON, C. PLAYOUST, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24**(1997), 235–265.
- [2] J. BOSMAN, P. BRUIN, A. DUJELLA, F. NAJMAN, *Ranks of elliptic curves with prescribed torsion over number fields*, preprint.
- [3] J. E. CREMONA, *Algorithms for Modular Elliptic Curves*, Cambridge Univ. Press, 1997.

- [4] T. DOKCHITSER, *Notes on the Parity Conjecture*, to appear in a CRM Advanced Courses volume, Birkhauser.
- [5] A. DUJELLA, M. JUKIĆ BOKUN, *On the rank of elliptic curves over  $\mathbb{Q}(i)$  with torsion group  $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$* , Proc. Japan Acad. Ser. A Math. Sci. **86**(2010), 93–96.
- [6] M. JUKIĆ BOKUN, *On the rank of elliptic curves over  $\mathbb{Q}(\sqrt{-3})$  with torsion groups  $\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$  and  $\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$* , Proc. Japan Acad. Ser. A Math. Sci. **87**(2011), 61–64.
- [7] S. KAMIENNY, *Torsion points on elliptic curves and  $q$ -coefficients of modular forms*, Invent. Math. **109**(1992), 221–229.
- [8] S. KAMIENNY, F. NAJMAN, *Torsion groups of elliptic curves over quadratic fields*, Acta. Arith. to appear.
- [9] M. A. KENKU, F. MOMOSE, *Torsion points on elliptic curves defined over quadratic fields*, Nagoya Math. J. **109**(1988), 125–149.
- [10] L. KULESZ, *Families of elliptic curves of high rank with nontrivial torsion group over  $\mathbb{Q}$* , Acta Arith. **108**(2003), 339–356.
- [11] B. MAZUR, *Rational isogenies of prime degree*, Invent. Math. **44**(1978), 129–162.
- [12] F. P. RABARISON, *Structure de torsion des courbes elliptiques sur les corps quadratiques*, Acta Arith. **144**(2010), 17–52.
- [13] D. SIMON, *Le fichier gp*, <http://www.math.unicaen.fr/~simon/ell.gp>.

Mirela Jukić Bokun  
 Department of Mathematics, University of Osijek,  
 Trg Ljudevita Gaja 6, 31000 Osijek, Croatia  
*E-mail address:* mirela@mathos.hr