

# On the $l_p$ -norm estimation in a quasilinear regression model

DRAGAN JUKIĆ

*Department of Mathematics, J.J. Strossmayer University of Osijek, Trg  
Ljudevita Gaja 6, HR-31 000 Osijek, Croatia*

**Abstract.** This talk will be based on my recently submitted manuscript on the  $l_p$ -norm ( $1 \leq p < \infty$ ) estimation of the parameters in a quasilinear regression model of the form

$$g(t; \boldsymbol{\alpha}) = \varphi(f_0(t) + \alpha_1 f_1(t) + \cdots + \alpha_n f_n(t)),$$

where  $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_n)^T \in \mathbb{R}^n$  is an unknown vector parameter,  $f_0, f_1, \dots, f_n$  are arbitrary fixed functions, and the function  $\varphi : I \rightarrow \mathbb{R}$ , with  $I \subseteq \mathbb{R}$  being an interval (open, closed, half-open, bounded or unbounded), is continuous and strictly monotonic.

Many important model functions which often appear in applied research are quasilinear or can be parameterized as a quasilinear model. For example: When  $\varphi(u) = \exp(u)$ , we have exponential regression; when  $\varphi(u) = u^a$ , where  $a \neq 0$  is given, we have power regression; when  $\varphi(u) = 1/u$ , we have hyperbolic regression.

The focus of this talk will be on the existence of the best  $l_p$ -norm estimator in a quasilinear regression model of the above form. I will review what is known about this problem and then present a theorem which guarantees the existence of the best  $l_p$ -norm estimator. From that theorem, which both extends and generalizes the previously known existence result, the existence of the best  $l_p$ -norm estimator for the whole class of nonlinear model functions follows immediately.