

## An invariance principle for the law of the iterated logarithm for vector-valued additive functionals of Markov chains

GUANGYU YANG<sup>1</sup>, YU MIAO<sup>2,\*</sup> AND XIAOCAI ZHANG<sup>3</sup>

<sup>1</sup> *Department of Mathematics, Zhengzhou University, Zhengzhou, 450 001, P. R. China*

<sup>2</sup> *College of Mathematics and Information Science, Henan Normal University, Xinxiang, 453 007, P. R. China*

<sup>3</sup> *College of Science, Henan University of Technology, Zhengzhou, 450 001, P. R. China*

Received April 24, 2011; accepted October 15, 2012

---

**Abstract.** In this note, we prove the Strassen's strong invariance principle for vector-valued additive functionals of a Markov chain via the martingale argument and the theory of fractional coboundaries.

**AMS subject classifications:** 60F05, 60J05

**Key words:** additive functionals of Markov chains, law of the iterated logarithm, Dunford-Schwarz operator, fractional coboundaries, vector-valued martingales

---

### 1. Introduction

Let  $(X_n)_{n \geq 0}$  denote a stationary ergodic Markov chain defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ , with values in a measurable space  $(\mathcal{X}, \mathcal{B})$ . Let  $Q(x, dy)$  be its transition kernel and  $\pi$  the stationary initial distribution. Fix an integer  $d \geq 1$  and for  $p \geq 1$  let  $L^p(\pi)$  denote the space of (equivalence classes of)  $\mathcal{B}$ -measurable functions  $g : \mathcal{X} \rightarrow \mathbb{R}^d$  such that  $\|g\|_p^p := \int_{\mathcal{X}} |g(x)|^p \pi(dx) < \infty$ , and let  $L_0^p(\pi)$  denote the set of  $g \in L^p(\pi)$  for which  $\int_{\mathcal{X}} g d\pi = 0$ . Here,  $|\cdot|$  denotes the Euclidean norm on  $\mathbb{R}^d$ .

Now fix an  $\mathbb{R}^d$ -valued function  $g \in L_0^2(\pi)$ . For  $n \geq 0$ , define

$$S_{n+1} = S_{n+1}(g) := \sum_{i=0}^n g(X_i) \quad \text{and} \quad S_0 = 0. \quad (1)$$

For the question of central limit type results for  $S_n$ , there have been numerous studies from many angles and under different assumptions; see Maxwell and Woodroffe [6], Derriennic and Lin [4] and references therein.

This short note is a natural continuation of Maxwell and Woodroffe [6]. Our goal is to consider the problem that  $S_n$  satisfies the law of the iterated logarithm (LIL) under some proper conditions. Since the appearance of Strassen's paper [13], almost sure invariance principles for the law of iterated logarithm have been obtained for a large class of independent and dependent sequence  $(Y_n)_{n \geq 1}$ ; see Hall and Heyde [5], and Philipp and Stout [10]. Here, the Skorokhod representation plays an important

---

\*Corresponding author. *Email addresses:* guangyu@zzu.edu.cn (G. Yang), yumiao@htu.cn (Y. Miao), xc.zh@haut.edu.cn (X. Zhang)



role. However, we encounter the essential difficulties, when considering the vector-valued martingale, since Monrad and Philipp [9] proved that it is impossible to embed a general  $\mathbb{R}^d$ -valued martingale in an  $\mathbb{R}^d$ -valued Gaussian process.

In this paper, we mainly take along the lines of Maxwell and Woodroffe [6], and use Berger's strong approximation [1]. Moreover, we identify the limsup in the functional LIL just the square root of the trace of the diffusion matrix corresponding to the functional central limit theorems (CLT). Recently, Zhao and Woodroffe [14] published a LIL for stationary processes that is stronger than our results in the case of  $d = 1$ , please refer to Miao and Yang [8]. However, the present note has the advantage to make clear that the reasonings used to get the functional CLT in its "almost sure" sense lead easily to the functional form of the multidimensional LIL (Zhao and Woodroffe [14] considered only the classical scalar form of this law but their reasoning could apply to a more general situation).

## 2. Main results

For introducing our main results, we need some notations. Let  $C([0, 1], \mathbb{R}^d)$  be the Banach space of continuous maps from  $[0, 1]$  to  $\mathbb{R}^d$ , endowed with the supremum norm  $\|\cdot\|$ , using the Euclidean norm in  $\mathbb{R}^d$ . Denote by  $K$  the set of absolutely continuous maps  $f \in C([0, 1], \mathbb{R}^d)$ , such that

$$f(0) = 0, \quad \int_0^1 |\dot{f}(t)|^2 dt \leq 1,$$

where  $\dot{f}$  denotes the derivative of  $f$  determined almost everywhere with respect to Lebesgue measure. Obviously,  $K$  is relatively compact and closed. Define

$$\xi_n(t) = (2n \log \log n)^{-1/2} [S_k + (nt - k)g(X_k)]$$

for  $t \in [\frac{k}{n}, \frac{k+1}{n})$ ,  $k = 0, 1, 2, \dots, n-1$ . In order to avoid difficulties in specification, we adopt the convention that  $\log \log x = 1$ , if  $0 < x \leq e^e$ . Then,  $\xi_n$  is a random element with values in  $C([0, 1], \mathbb{R}^d)$ . In addition, given a function  $h \in L^1(\pi)$ , we define an operator

$$Qh(x) = \int h(y)Q(x, dy), \quad \pi\text{-a.e. } x \in \mathcal{X}.$$

Obviously,  $Q$  is a contraction on  $L^p(\pi)$  for  $p \geq 1$ .

**Theorem 1.** *Let  $g \in L_0^2(\pi)$  and assume that there exists an  $\alpha \in (0, 1/2)$  such that*

$$\left\| \sum_{i=0}^{n-1} Q^i g \right\|_2 = O(n^\alpha). \quad (2)$$

*Then, the sequence of functions  $(\xi_n(\cdot), n \geq 1)$  is almost surely relatively compact in the space  $C([0, 1], \mathbb{R}^d)$ , and the set of its limit points as  $n \rightarrow \infty$ , coincides with  $\sqrt{\text{tr}(\mathfrak{D})}K$ , where  $\text{tr}(\cdot)$  denotes the trace operator of a matrix and  $\mathfrak{D} = \mathbb{E}(M_1 M_1^t) = \int H H^t d\pi_1$  is the diffusion matrix corresponding to the functional central limit theorem, where  $M_1$ ,  $H$  and  $\pi_1$  are defined in Section 3.*



**Theorem 2.** *Let  $g \in L_0^2(\pi)$  and assume that there exists an  $\alpha \in (0, 1/2)$  for which (2) is satisfied. Then*

$$\limsup |S_n|/\sqrt{2n \log \log n} = \sqrt{\text{tr}(\mathfrak{D})}, \quad \mathbb{P}\text{-a.s.} \quad (3)$$

It is worthwhile to give some comments on Theorem 1 and Theorem 2.

**Remark 1.** *Maxwell and Woodroffe [6] showed the functional CLT for  $S_n$  when  $d = 1$ , under condition (2). However, the CLT is proved there under the following weaker condition*

$$\sum_{n=1}^{\infty} n^{-3/2} \left\| \sum_{i=0}^{n-1} Q^i g \right\|_2 < \infty. \quad (4)$$

Recently, for  $d = 1$ , Peligrad and Utev [11] proved the functional CLT under condition (4) by developing some new maximal inequality for stationary sequences. It is well known that the LIL is closely related to the CLT in some sense, hence the functional LIL should be true under condition (4). But, under our present framework, the functional LIL cannot be proved under condition (4) since the fractional coboundary theory of Derriennic and Lin [3] is no longer valid. Cuny [2] also treated the LIL for  $S_n$  when  $d = 1$  under some conditions. For an update account of condition (4) and some other forms, please refer to Merlevède, Peligrad and Peligrad [7] and references therein.

### 3. Proof of main results

For  $\varepsilon > 0$ , let  $h_\varepsilon$  be the solution of the equation

$$(1 + \varepsilon)h = Qh + g,$$

where  $Q$  is defined as in Section 2. In fact,

$$h_\varepsilon = \sum_{n=1}^{\infty} (1 + \varepsilon)^{-n} Q^{n-1} g. \quad (5)$$

Note that  $h_\varepsilon \in L^p(\pi)$ , if  $g \in L^p(\pi)$ . Let  $\pi_1$  be the joint distribution of  $X_0$  and  $X_1$ , so that  $\pi_1(dx_0, dx_1) = Q(x_0, dx_1)\pi(dx_0)$ ; denote the  $L^2$ -norm on  $L^2(\pi_1)$  by  $\|\cdot\|_1$ ; and let

$$H_\varepsilon(x_0, x_1) = h_\varepsilon(x_1) - Qh_\varepsilon(x_0)$$

for  $x_0, x_1 \in \mathcal{X}$ . For any  $\varepsilon > 0$ , let

$$M_n(\varepsilon) = \sum_{i=0}^{n-1} H_\varepsilon(X_i, X_{i+1}) \quad \text{and} \quad R_n(\varepsilon) = Qh_\varepsilon(X_0) - Qh_\varepsilon(X_n),$$

hence, by simple computations,

$$S_n(g) = M_n(\varepsilon) + \varepsilon S_n(h_\varepsilon) + R_n(\varepsilon). \quad (6)$$

For convenience, we summarize the results of Maxwell and Woodroffe [6] as the following theorem.



**Theorem 3** (Theorem MW). *Assume that  $g \in L_0^2(\pi)$  and that there exists an  $\alpha \in (0, 1/2)$  for which (2) is satisfied. Then we have*

1. *The limit  $H = \lim_{\varepsilon \rightarrow 0^+} H_\varepsilon$  exists in  $L^2(\pi_1)$ . Moreover, if one defines*

$$M_n = \sum_{i=0}^{n-1} m_i,$$

*where  $m_i = H(X_i, X_{i+1})$ , then  $(m_n)_{n \geq 0}$  is a stationary and ergodic  $\mathbb{P}$ -square integrable martingale difference sequence, with respect to the filtration  $\{\mathcal{F}_n = \sigma(X_0, \dots, X_n)\}_{n \geq 0}$ ;*

2.  *$\|h_\varepsilon\|_2 = O(\varepsilon^{-\alpha})$ , and if  $R_n = S_n - M_n = M_n(\varepsilon) - M_n + \varepsilon S_n(h_\varepsilon) + R_n(\varepsilon)$ , then*

$$\mathbb{E}(|R_n|^2) = O(n^{2\alpha}).$$

### 3.1. Proof of Theorem 1

For  $0 \leq t \leq 1$ , define

$$\begin{aligned} \zeta_n(t) &= (2n \log \log n)^{-1/2} M_{[nt]}, \\ \eta_n(t) &= (2n \log \log n)^{-1/2} B(nt), \end{aligned}$$

where  $M_n$  is as defined in Section 2 and  $B(\cdot)$  is an  $\mathbb{R}^d$ -valued Brownian motion with mean 0 and diffusion matrix  $\mathfrak{D}$ . Theorem 1 of Strassen [13] shows that  $(\eta_n(\cdot))_{n \geq 1}$  is relatively compact and the set of its limit points coincides with  $\sqrt{\text{tr}(\mathfrak{D})}K$ .

Notice that by part (1) of Theorem MW,  $(M_n)_{n \geq 1}$  is a square integrable martingale with strictly stationary increments. Moreover,

$$\mathbb{E}(\langle u, m_0 \rangle^2) < \infty \quad \text{and} \quad \mathbb{E}(\langle u, m_0 \rangle) = 0, \quad \text{for all } u \in \mathbb{R}^d, \quad (7)$$

where,  $\langle \cdot, \cdot \rangle$  denotes the inner product in  $\mathbb{R}^d$ . Therefore, Corollary 4.1 of Berger [1] implies that,

*Without changing its distribution, one can redefine the sequence  $(M_n)_{n \geq 1}$  on a new probability space  $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{\mathbb{P}})$  on which there exists an  $\mathbb{R}^d$ -valued Brownian motion  $(B(t))_{t \geq 0}$  with mean 0 and diffusion matrix  $\mathfrak{D}$  such that*

$$|M_{[t]} - B(t)| = o((t \log \log t)^{1/2}), \quad \hat{\mathbb{P}}\text{-a.s.} \quad (\text{as } t \rightarrow \infty). \quad (8)$$

where,  $\mathfrak{D} = \lim_{n \rightarrow \infty} n^{-1} \text{Cov}(M_n)$ .

**Remark 2.** *Birkhoff-Khinchin's ergodic theorem and a simple calculation show that  $\mathfrak{D} = \mathbb{E}(M_1 M_1^t) = \int H H^t d\pi_1$  is the diffusion matrix corresponding to the functional central limit theorem; please see Rassoul-Agha and Seppäläinen [12].*

That is to say,

$$\sup_{0 \leq t \leq 1} |M_{[nt]} - B(nt)| = o((2n \log \log n)^{1/2}), \quad \mathbb{P}\text{-a.s.}$$



Hence,

$$\begin{aligned} ||\zeta_n - \eta_n|| &= (2n \log \log n)^{-1/2} \sup_{0 \leq t \leq 1} |M_{[nt]} - B(nt)| \\ &= o(1), \quad \mathbb{P}\text{-a.s.} \end{aligned}$$

Define

$$\tilde{\zeta}_n(t) = (2n \log \log n)^{-1/2} [M_k + (nt - k)m_k]$$

for  $t \in [\frac{k}{n}, \frac{k+1}{n})$ ,  $k = 0, 1, 2, \dots, n-1$ . Then  $\tilde{\zeta}_n \in C([0, 1], \mathbb{R}^d)$  and

$$\sup_{t \in [0, 1]} |\zeta_n(t) - \tilde{\zeta}_n(t)| = (2n \log \log n)^{-1/2} \max_{0 \leq k \leq n-1} |m_k|.$$

Next, we give the order estimation of  $\max_{0 \leq k \leq n-1} |m_k|$ . Since the fact that

$$\frac{1}{n} \sum_{k=1}^n |m_k|^2 \longrightarrow \text{tr}(\mathfrak{D}), \quad \mathbb{P}\text{-a.s.}$$

we have  $\max_{0 \leq k \leq n-1} |m_k| = o(n^{1/2})$  a.s. Hence,

$$||\zeta_n - \tilde{\zeta}_n|| = \sup_{t \in [0, 1]} |\zeta_n(t) - \tilde{\zeta}_n(t)| = o(1), \quad \mathbb{P}\text{-a.s.}$$

The above discussions immediately yield the following claim:

$(\tilde{\zeta}_n(\cdot), n \geq 1)$  is almost surely relatively compact and the set of its limit points coincides with  $\sqrt{\text{tr}(\mathfrak{D})}K$ .

We now turn to deal with the negligible term  $R_n$  in the sense of functional LIL. Firstly, let us recall the concept of the Dunford-Schwarz (DS) operator; see Derriennic and Lin [3]. We call  $T$  a DS operator on  $L^1$  of a probability space, if  $T$  is a contraction of  $L^1$  such that  $\|Tf\|_\infty \leq \|f\|_\infty$  for every  $f \in L^\infty$ . If  $\theta$  is a measure preserving transformation in a probability space  $(\Omega, \Sigma, \mu)$ , then the operator  $Tf = f \circ \theta$  is a DS operator on  $L^1(\mu)$ . More generally, any Markov transition operator  $P$  with an invariant probability measure yields a positive DS operator.

**Lemma 1** (Lemma DL, [3]).

1. Let  $T$  be a contraction in a Banach space  $X$ , and let  $0 < \beta < 1$ . If

$$\sup_n \left\| \frac{1}{n^{1-\beta}} \sum_{k=1}^n T^k y \right\| < \infty, \text{ then } y \in (I - T)^\alpha X \text{ for every } 0 < \alpha < \beta.$$

2. Let  $T$  be a DS operator in  $L^1(\mu)$  of a probability space, and fix  $1 < p < \infty$ , with  $q = p/(p-1)$ . Let  $0 < \alpha < 1$ , and  $f \in (I - T)^\alpha L^p$ . If  $\alpha > 1 - \frac{1}{p} = \frac{1}{q}$ ,

$$\text{then } \frac{1}{n^{1/p}} \sum_{k=0}^{n-1} T^k f \rightarrow 0 \text{ a.e.}$$



Note that Lemma DL was originally proved for  $d = 1$ ; however, we can make the shift operating coordinate-wise, hence the original proof of Derriennic and Lin is enough for our case. To apply the above lemma, i.e., the theory of fractional coboundaries named by Derriennic and Lin [3], we need to construct a DS operator. On  $\mathcal{X} \times \mathcal{X}$ , define

$$f(x_0, x_1) = g(x_0) - H(x_0, x_1),$$

then

$$\begin{aligned} R_n &= S_n - M_n \\ &= \sum_{i=0}^{n-1} [g(X_i) - H(X_i, X_{i+1})] \\ &= \sum_{i=0}^{n-1} f(X_i, X_{i+1}). \end{aligned} \tag{9}$$

Let  $\theta$  be the shift map on the path space  $\mathcal{X}^{\mathbb{N}}$  for the Markov chain which is a contraction on  $L^2(\mathbb{P})$ . The DS operator is the shift  $\theta$ . For a sequence  $x = (x_i)_{i \in \mathbb{N}} \in \mathcal{X}^{\mathbb{N}}$ , define  $F(x) = f(x_0, x_1)$ , then we have

$$F \in L^2(\mathbb{P}) \quad \text{and} \quad R_n = \sum_{k=0}^{n-1} F \circ \theta^k.$$

From part (2) of Theorem MW, there exists a constant  $1/2 < \beta < 1 - \alpha$ , such that

$$\sup_n \left\| \frac{1}{n^{1-\beta}} \sum_{k=0}^{n-1} F \circ \theta^k \right\| < \infty, \tag{10}$$

Since part (1) of Lemma DL and  $0 < \alpha < 1/2$ , we have  $F \in (I - \theta)^\eta L^2(\mathbb{P})$ , for some  $\eta \in (1/2, 1 - \alpha)$ . By part (2) of Lemma DL, we have

$$\frac{1}{n^{1/2}} R_n \rightarrow 0, \quad \mathbb{P}\text{-a.s.}$$

Furthermore, applying an elementary property of real convergent sequences, we immediately get

$$\max_{0 \leq k \leq n} |R_k| = o((2n \log \log n)^{1/2}), \quad \mathbb{P}\text{-a.s.}$$

Consequently,

$$(2n \log \log n)^{-1/2} \sup_{0 \leq t \leq 1} |R_{[nt]}| \rightarrow 0, \quad \mathbb{P}\text{-a.s.} \tag{11}$$

From the above discussions, we complete the proof of Theorem 1.



### 3.2. Proof of Theorem 2

Here, we take along the lines of the proof of Theorem 4.8 in Hall and Heyde [5]. Let  $\{e_i\}_{i=1}^d$  the canonical basis of  $\mathbb{R}^d$ . For any  $\mathbb{R}^d$ -valued function  $f$ , denote  $f = (f_1, f_2, \dots, f_d)^t$ . By the definition of  $K$ , for any  $f \in \sqrt{\text{tr}(\mathfrak{D})}K$ , we have

$$|f(t)|^2 = \sum_{i=1}^d \left( \int_0^t \dot{f}_i(s) ds \right)^2 \leq \sum_{i=1}^d \left( \int_0^t \dot{f}_i(s)^2 ds \right) \int_0^t 1 ds \leq \text{tr}(\mathfrak{D})t \quad (12)$$

where the first inequality is due to the Cauchy-Schwarz's inequality. So,  $|f(t)| \leq \sqrt{\text{tr}(\mathfrak{D})t}$ . It follows that  $\sup_{t \in [0,1]} |f(t)| \leq \sqrt{\text{tr}(\mathfrak{D})}$ . Hence, by Theorem 1

$$\limsup_n \sup_{t \in [0,1]} |\xi_n(t)| \leq \sqrt{\text{tr}(\mathfrak{D})}, \quad \mathbb{P} - a.s. \quad (13)$$

and setting  $t = 1$ ,

$$\limsup_n |S_n| / \sqrt{2n \log \log n} \leq \sqrt{\text{tr}(\mathfrak{D})}, \quad \mathbb{P} - a.s. \quad (14)$$

On the other hand, we put  $f(t) = t \sqrt{\frac{\text{tr}(\mathfrak{D})}{d}} \sum_{i=1}^d e_i$ ,  $t \in [0, 1]$ . Then,  $f \in \sqrt{\text{tr}(\mathfrak{D})}K$  and so for  $\mathbb{P} - a.s. \omega$ , there exists a sequence  $n_k = n_k(\omega)$  such that

$$\xi_{n_k}(\cdot)(\omega) \xrightarrow{\|\cdot\|} f(\cdot). \quad (15)$$

Particularly,  $f(1) = \sqrt{\frac{\text{tr}(\mathfrak{D})}{d}} \sum_{i=1}^d e_i$ ,  $|\xi_{n_k}(1)(\omega)| \rightarrow |f(1)|$ . That is to say,

$$|S_{n_k}| / \sqrt{2n_k \log \log n_k} \rightarrow \sqrt{\text{tr}(\mathfrak{D})}, \quad \mathbb{P} - a.s. \quad (16)$$

This completes the proof of Theorem 2.

### Acknowledgment

The authors wish to thank Prof. Y. Derriennic and Prof. M. Lin for sending the key reference paper [3] and the anonymous reviewers for their critical comments which improve the presentation of this note. The work of G. Yang was supported by NSFC (11201431) and Foundations of Zhengzhou University and that of Y. Miao was supported by NSFC (11001077), NCET (NCET-11-0945), Henan Province Foundation and Frontier Technology Research Plan (112300410205) and Plan For Scientific Innovation Talent of Henan Province (124100510014).

### References

- [1] E. BERGER, *An almost sure invariance principle for stationary ergodic sequences of Banach space valued random variables*, Probab. Th. Rel. Fields. **84**(1990), 161–201.
- [2] C. CUNY, *Pointwise ergodic theorems with rate and application to limit theorems for stationary processes*, Stoch. Dynam **11**(2011), 135–155.



- [3] Y. DERRIENNIC, M. LIN, *Fractional Poisson equations and ergodic theorems for fractional coboundaries*, Israel J. Math. **123**(2001), 93–130.
- [4] Y. DERRIENNIC, M. LIN, *The central limit theorem for Markov chains started at a point*, Probab. Th. Rel. Fields. **125**(2003), 73–76.
- [5] P. HALL, C. C. HEYDE, *Martingale limit theory and its applications*, Academic Press, New York, 1980.
- [6] M. MAXWELL, M. WOODROOFE, *Central limit theorems for additive functionals of Markov chains*, Ann. Probab. **28**(2000), 713–724.
- [7] F. MERLEVÈDE, C. PELIGRAD, M. PELIGRAD, *Almost sure invariance principles via martingale approximation*, Stoch. Process. Appl. **122**(2012), 170–190.
- [8] Y. MIAO, G. Y. YANG, *The law of the iterated logarithm for additive functionals of Markov chains*, Statist. Probab. Letters. **78**(2008), 265–270.
- [9] D. MONRAD, W. PHILIPP, *The problem of embedding vector-valued martingales in Gaussian process*, Teor. Veroyatn. Primen. **35**(1990), 384–387.
- [10] W. PHILIPP, W. F. STOUT, *Almost sure invariance principles for partial sums of weakly dependent random variables*, Memor. Amer. Math. Soci. No. 161, Amer. Math. Soc., Providence 1975.
- [11] M. PELIGRAD, S. UTEV, *A new maximal inequality and invariance principle for stationary sequences*, Ann. Probab. **38**(2005), 798–815.
- [12] F. RASSOUL-AGHA, T. SEPPÄLÄINEN, *An almost surely invariance principle for additive functionals of Markov chains*, Statist. Probab. Letters. **78**(2008), 854–860.
- [13] V. STRASSEN, *An invariance principle for the law of the iterated logarithm*, Z. Wahrs. verw. Geb. **3**(1964), 211–226.
- [14] O. ZHAO, M. WOODROOFE, *Law of the iterated logarithm for stationary processes*, Ann. Probab. **36**(2008), 127–142.